

学校编码: 10384

分类号_____密级_____

学 号: 20620101151424

UDC_____

厦 门 大 学

硕 士 学 位 论 文

一氧化碳低温氧化负载型钯催化剂的侧柏
叶浸出液辅助还原制备与性能研究

Study on Supported Pd Catalysts Biosynthesized by *Cacumen
Platycladi* Extract and their Catalytic Performance on Low
Temperature Oxidation of CO

国家自然科学基金重点项目(21036004)资助

朱 婧

指导教师姓名: 李清彪 教 授

孙道华 副教授

专 业 名 称: 工 业 催 化

论文提交日期: 2 0 1 3 年 月

论文答辩日期: 2 0 1 3 年 月

学位授予日期: 2 0 1 3 年 月

答辩委员会主席: _____

评 阅 人: _____

2013 年 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

（ ） 1.经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。

（ ） 2.不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月

摘 要

CO 低温氧化反应在汽车尾气处理、封闭式-循环 CO₂ 激光器、氢燃料电池中少量 CO 的消除及作为催化领域模型反应等方面都有重要应用。近年来, 研究者致力于改进催化剂的制备方法, 以期获得低温下高活性、高稳定性的催化剂。本论文将植物还原法引入 Pd 催化剂的制备过程, 分别采用溶胶负载法 (SI 法) 和吸附还原法 (AR 法) 制备负载型 Pd 催化剂, 将所制备的 Pd 催化剂进行表征, 并用于 CO 低温氧化体系中, 考察催化剂制备条件对反应性能的影响, 优化催化剂的制备条件, 探索催化剂的结构和表面性质与催化反应性能之间的关系。

首先采用 SI 法制备了 Pd 催化剂, 研究结果表明: 通过改变反应温度、侧柏叶浸出液浓度及 Pd 前驱体的浓度可以调节 Pd 纳米颗粒的粒径, Pd 颗粒粒径越小催化活性越高。侧柏叶浸出液浓度过高, 虽然会获得粒径较小的 Pd 纳米颗粒, 但同时也覆盖了 Pd 催化剂的活性位点, 导致催化剂活性降低。经过焙烧处理去除催化剂表面的生物物质, 催化剂活性提高, 此时催化剂表面为 Pd²⁺物种, 说明氧化态的 Pd²⁺可作为该反应的活性物种。经过氢气还原处理后, 催化剂表面部分 Pd²⁺被还原为 Pd⁰, 导致催化活性再次提升, 表明 Pd⁰物种也是 CO 氧化体系中催化剂表面的活性物种。催化剂活性数据结果表明: 溶胶负载法制备, 以 Pd(NO₃)₂ 为前驱体, CeO₂ 为载体, Pd 理论负载量为 0.5 wt.%, 侧柏叶浸出液浓度为 5 g/L, 空气气氛下 300 °C 焙烧 2 h, 氢气中 500 °C 还原处理 2 h 制得的 Pd 催化剂活性较优。在空速 6000 mL/g_{cat}/h, 原料气比例 CO/O₂/N₂ = 1/1/98, 反应温度 30 °C 时 CO 转化率为 49.6 %, 50 °C 时 CO 转化率可达到 100 %。

采用 AR 法制备了 Pd 催化剂, 实验结果表明: 以不含 Cl⁻离子的 Pd(NO₃)₂ 为前驱体可以获得较高的催化活性, 以 CeO₂ 为载体, 以侧柏叶浸出液做还原剂制备所得的催化剂活性较优。该法不但可以有效地将 Pd 负载于 CeO₂ 载体上, 而且在高负载情况下亦可以实现 Pd 的高分散。和 SI 法所获催化剂类似, 在空气氛围中一定温度下焙烧催化剂可以除去催化剂表面残留的生物物质, 暴露出催化剂的活性位点从而提高催化剂的活性, 同样经过焙烧之后的催化剂表面 Pd 物种被氧化为 Pd²⁺, 此时的 Pd²⁺不是以 PdO 形式而是以 Pd_xCeO_{2-δ} 固溶体的形式存在。对 AR 法制备的 Pd/CeO₂ 进行氢气还原处理, 还原后催化剂活性有了极大的提高。对催化剂进行稳定性评价, 发现催化剂活性下降后, 再次通氢处理可以恢复催化

剂的活性，并且可以延长催化剂的寿命。这可能是因为反应过程中 Pd^0 被氧化，而通氢处理可以使 Pd 物种恢复到金属态 Pd^0 ，并加深了催化剂表面 Pd 的还原程度。另外本文还初步探索了 CeO_2 形貌对催化剂活性的影响，发现合成的纳米棒状和纳米方块状 CeO_2 负载 Pd 之后都可以获得比常规 CeO_2 载体更高的活性。催化剂活性数据显示：吸附还原法制备，以 $\text{Pd}(\text{NO}_3)_2$ 为前驱体，纳米棒状 CeO_2 为载体，Pd 理论负载量为 0.5 wt.%，侧柏叶浸出液浓度为 5 g/L，空气气氛下 300 °C 焙烧 2 h，氢气中 300 °C 还原处理 2 h 制得的 Pd 催化剂活性较优。在空速 6000 mL/g_{cat}/h，原料气比例 $\text{CO}/\text{O}_2/\text{N}_2 = 1/1/98$ ，反应温度 30 °C 时 CO 转化率达 62 %，50 °C 时 CO 转化率可达 100 %。

关键字： 负载型 Pd 催化剂；CO 低温氧化；生物还原；二氧化铈

Abstract

The catalytic oxidation of CO at low temperature has attracted considerable attention because of its wide applications in automotive emission control, exhaust abatement for CO₂ lasers, CO preferential oxidation for proton exchange membrane fuel cells and as a prototypical reaction for heterogeneous processes. It is necessary to introduce eco-friendly and low-priced process to prepare effective catalysts with high activity and stability for CO oxidation at low temperature. In this paper, a new green method integrating bio-reduction and immobilization was developed to prepare supported Pd catalysts for CO oxidation. Specifically, supported Pd catalysts were prepared by sol-immobilization method (SI) and adsorption-reduction (AR) method respectively using plant extract as reductant and stabilizing agent. The influence of catalysts preparation conditions on CO catalytic activity were investigated and the catalysts were characterized by a variety of techniques to research the relationship between structure and surface properties of catalysts and their catalytic performance.

Firstly, SI method was employed to prepare supported Pd catalysts. The results showed that the size of Pd nanoparticles (PdNPs), which had significant impact on catalytic performance, could be controlled by the adjustment of reaction temperature, concentration of *Cacumen Platycladi* (CP) extract and Pd precursor. And smaller PdNPs had a better catalytic activity in CO oxidation. However, smaller PdNPs achieved with a high CP concentration caused a low activity because the surface active sites were covered by the biomass. Calcination could remove the biomass residual and expose the active size on supported Pd catalysts, accordingly the catalytic activity of catalysts increased after calcination. The Pd species on the catalysts surface were in Pd²⁺ state, which showed that Pd²⁺ could act in CO oxidation. It was also found that H₂ could remarkably improve the conversion of CO oxidation at low temperature. After H₂ reduction, Pd²⁺ on the catalysts surface turned to Pd⁰, which proved the Pd⁰ was more effective and active species than Pd²⁺ in CO oxidation at low temperature. As a result, the optimum conditions of preparing Pd/CeO₂ catalyst was as follows: Pd loading of 0.5 wt.%, CP extract concentration of 5 g/L, calcination at 300

°C for 2 h in air, hydrogen reduction at 500 °C for 2 h. In this condition, CO conversion reached 49.6 % at 30 °C and 100 % at 50 °C under a 6000 mL/g_{cat}/h space velocity with CO/O₂/N₂ = 1/1/98.

Secondly, AR method was then used in the biosynthesis of supported Pd catalysts. The results revealed that the kinds of Pd precursors, supports, plants were highly affecting catalytic performance. It was found Pd(NO₃)₂ as precursor, CeO₂ as support and CP extract as biomass reductant could achieve high CO conversion. AR method could not only effectively load Pd on CeO₂, but also achieve high dispersion of Pd in the case of high Pd loading amount. Similar to SI method, calcination removed the biomass residual to expose the active sites and improved the catalytic activity. Whereas, Pd species in the catalyst prepared by AR method were existed in Pd_xCeO_{2-δ} phase instead of PdO after calcinations in air. Also, H₂ reduction could significantly enhance the catalytic activity by partially reduce Pd²⁺ to Pd⁰ species, which as well confirmed that Pd⁰ was more active than Pd²⁺ in CO oxidation reaction. The catalyst stability was evaluated and it was found that not only the catalyst activity could recover but also the catalyst lifetime was prolonged after H₂-treatment. It might be because H₂ reduction treatment reduced the Pd²⁺ species formed in reaction process to more active Pd⁰. Furthermore, the influence of different ceria crystal morphology on the performance of catalysts was researched and the results showed that catalysts using nanocubes and nanorods CeO₂ as supports exhibit higher catalytic activity than that using conventional CeO₂. Hence, the optimum conditions to obtain Pd catalysts by AR method was found as follows: nanorods CeO₂ as support, the Pd loading of 0.5 wt%, CP extract concentration of 5 g/L, calcination at 300 °C for 2 h in air and then H₂ reduction at 300 °C for 2 h. In this condition, CO conversion reached 62 % at 30 °C and 100 % at 50 °C under a 6000 mL/g_{cat}/h space velocity with CO/O₂/N₂ = 1/1/98.

Keywords: Supported Pd catalyst; Low temperature CO oxidation; Bioreduction; Ceria

目 录

第一章 绪论	1
1.1 CO 低温氧化的研究现状及意义	1
1.2 CO 低温氧化催化剂	3
1.2.1 非贵金属氧化物催化剂	3
1.2.1.1 霍加拉特	3
1.2.1.2 钙钛矿	4
1.2.1.3 尖晶石	4
1.2.1.4 半导体	5
1.2.2 贵金属催化剂	6
1.2.2.1 Pt 催化剂	6
1.2.2.2 Au 催化剂	8
1.2.2.3 Pd 催化剂	11
1.3 负载型 Pd 催化剂的制备方法	13
1.3.1 化学法	13
1.3.1.1 浸渍法	13
1.3.1.2 共沉淀法	13
1.3.1.3 沉积-沉淀法	14
1.3.1.4 化学气相沉积法	14
1.3.2 生物还原法	15
1.3.2.1 微生物还原法	15
1.3.2.2 植物还原法	15
1.4 本论文的研究目标与研究内容	16
1.5 论文组成与概要	17
第二章 实验方法	19
2.1 主要原料和试剂	19
2.2 催化剂的制备	19

2.2.1 侧柏叶浸出液的制备	19
2.2.2 CeO ₂ 的制备	20
2.2.3 负载型 Pd 催化剂的制备	20
2.2.3.1 溶胶负载法	20
2.2.3.2 吸附还原法	20
2.3 催化剂的性能评价	21
2.4 催化剂的表征	22
2.4.1 X 射线粉末衍射	22
2.4.2 紫外-可见漫反射	22
2.4.3 透射电镜	22
2.4.4 热重分析	23
2.4.5 傅里叶变换红外光谱	23
2.4.6 X 射线光电子能谱	23
2.4.7 低温 N ₂ 物理吸附	23
2.4.8 原子吸收分光光谱	23
2.4.9 紫外可见吸收光谱	24
第三章 溶胶负载法制备 Pd 催化剂及其 CO 氧化性能考察	25
3.1 引言	25
3.2 Pd 溶胶的制备考察	25
3.2.1 温度的影响	25
3.2.2 侧柏叶浸出液浓度的影响	27
3.2.3 前驱体浓度的影响	28
3.2.4 FTIR 表征	29
3.3 催化剂制备条件的影响	30
3.3.1 载体的影响	30
3.3.1.1 TEM 表征	30
3.3.1.2 载体种类对催化活性的影响	31
3.3.2 制备温度的影响	32
3.3.2.1 XRD 表征	32

3.3.2.2 制备温度对催化剂活性的影响	33
3.3.3 侧柏叶浸出液浓度的影响	35
3.3.3.1 XRD 表征	35
3.3.3.2 侧柏叶浸出液浓度对催化剂活性的影响	36
3.3.4 焙烧的影响	37
3.3.4.1 XPS 表征	37
3.3.4.2 TG 表征	38
3.3.4.3 焙烧对催化活性的影响	39
3.3.5 前驱体对催化活性的影响	40
3.3.6 氢气还原的影响	41
3.3.6.1 XPS 表征	41
3.3.6.2 氢气还原对催化活性的影响	42
3.3.7 Pd 负载量对催化活性的影响	43
3.4 溶胶法制备 Au-Pd 催化剂初探	44
3.4.1 UV-Vis 表征	44
3.4.2 XRD 表征	46
3.4.3 TEM 表征	48
3.4.4 TG 表征	50
3.4.5 Au-Pd 比对催化活性的影响	50
3.5 本章小结	52
第四章 吸附还原法制备 Pd 催化剂及其 CO 氧化性能考察	54
4.1 引言	54
4.2 催化剂制备条件的影响	54
4.2.1 前驱体的影响	54
4.2.2 载体的影响	55
4.2.2.1 载体种类对催化活性的影响	55
4.2.2.2 XRD 表征	56
4.2.3 植物生物质种类的影响	57
4.2.3.1 植物生物质种类对催化活性的影响	57

4.2.3.2 TG 表征	58
4.2.4 侧柏叶浸出液浓度的影响	59
4.2.5 负载量的影响	60
4.2.5.1 XRD 表征	60
4.2.5.2 AAS 表征	61
4.2.5.3 低温 N ₂ 物理吸附表征	62
4.2.5.4 负载量对催化活性的影响	62
4.2.6 焙烧温度的影响	63
4.2.6.1 FTIR 表征和 TG 表征	63
4.2.6.2 XPS 表征	65
4.2.6.3 UV-Vis DRS 表征	65
4.2.6.4 TEM 表征	66
4.2.6.5 低温 N ₂ 物理吸附表征	67
4.2.6.6 焙烧温度对催化活性的影响	68
4.2.6.7 稳定性评价	70
4.2.7 氢气还原的影响	71
4.2.7.1 XRD 表征	71
4.2.7.2 XPS 表征	72
4.2.7.3 UV-Vis DRS 表征	73
4.2.7.4 氢气还原对催化活性的影响	74
4.2.8 CeO ₂ 载体的形貌的影响	75
4.2.8.1 TEM 表征	76
4.2.8.2 XRD 表征	76
4.2.8.3 低温 N ₂ 物理吸附表征	77
4.2.8.4 CeO ₂ 载体的形貌对催化活性的影响	77
4.2.9 催化剂 Pd/CeO ₂ /300/H ₂ 的稳定性评价	79
4.2.9.1 稳定性评价	79
4.2.9.2 XPS 表征	80
4.3 本章小结	81

第五章 结论与建议	83
5.1 结论	83
5.2 建议	84
参考文献	86
在学期间的科研成果	95
致 谢	96

Contents

Chapter 1 Introduction	1
1.1 The Background and Significance of the Reserch	1
1.2 Types of the catalysts for low-temperature CO oxidation	3
1.2.1 Base Metal Oxide Catalysts	3
1.2.1.1 Hopcalite	3
1.2.1.2 Perovskite.....	4
1.2.1.3 Spinel structure materials.....	4
1.2.1.4 Semiconductors.....	5
1.2.2 Noble Metal Catalysts.....	6
1.2.2.1 Supported Pt Catalysts	6
1.2.2.2 Supported Au Catalysts.....	8
1.2.2.3 Supported Pd Catalysts	11
1.3 Preparation method of Pd catalysts	13
1.3.1 Chemical method	13
1.3.1.1 Impregnation	13
1.3.1.2 Coprecipitation.....	13
1.3.1.3 Deposition-precipitation	14
1.3.1.4 Chemical vapour deposition	14
1.3.2 Biological method.....	15
1.3.2.1 Microorganism bioreduction method.....	15
1.3.2.2 Plant bioreduction method.....	15
1.4 Scope and objectives of this work.....	16
1.5 The Outline of the Thesis	17
Chapter 2 Experimental.....	19
2.1 Materials and reagents	19
2.2 Catalyst preparation.....	19
2.2.1 Preparation of CP extract.....	19
2.2.2 Preparation of CeO ₂	20

2.2.3	Preparation of Pd Catalysts	20
2.2.3.1	Sol-immobilization (SI) method	20
2.2.3.2	Adsorption–reduction (AR) method	20
2.3	Evaluation of catalytic properties.....	21
2.4	Catalyst characterization	22
2.4.1	XRD analysis	22
2.4.2	UV-Vis DRS analysis	22
2.4.3	TEM analysis	22
2.4.4	TG analysis	23
2.4.5	FTIR analysis	23
2.4.6	XPS analysis	23
2.4.7	BET analysis	23
2.4.8	AAS analysis.....	23
2.4.8	UV-Vis analysis	24
Chapter 3	Preparation of Pd catalysts by SI mehtod and their catalytic performance for CO oxidation.....	25
3.1	Introduction.....	25
3.2	Preparation of Pd sol	25
3.2.1	Effect of temperature	25
3.2.2	Effect of CP extract concentration	27
3.2.3	Effect of Pd precursor concentration	28
3.2.4	FTIR analysis	29
3.3	Effect of preparation conditions of catalysts	30
3.3.1	Effect of support	30
3.3.1.1	TEM analysis	30
3.3.1.2	Catalytic performance	31
3.3.2	Effect of preparation temperature	32
3.3.2.1	XRD analysis	32
3.3.2.2	Catalytic performance	33

3.3.3	Effect of CP extract concentration	35
3.3.3.1	XRD analysis	35
3.3.3.2	Catalytic performance	36
3.3.4	Effect of calcination	37
3.3.4.1	XPS analysis	37
3.3.4.2	TG analysis	38
3.3.4.3	Catalytic performance	39
3.3.5	Effect of Pd precursor	40
3.3.6	Effect of H ₂ -treatment	41
3.3.6.1	XPS analysis	41
3.3.6.2	Catalytic performance	42
3.3.7	Effect of Pd loading	43
3.4	Tentative exploration of Au-Pd catalysts	44
3.4.1	UV-Vis analysis	44
3.4.2	XRD analysis	46
3.4.3	TEM analysis	48
3.4.4	TG analysis	50
3.4.5	Effect of Au-Pd mass ratio	50
3.5	Summary	52
Chapter 4 Preparation of Pd catalysts by AR method and their catalytic performance for CO oxidation		54
4.1	Introduction	54
4.2	Effect of preparation conditions of catalysts	54
4.2.1	Effect of Pd precursor	54
4.2.2	Effect of support	55
4.2.2.1	Catalytic performance	55
4.2.2.2	XRD analysis	56
4.2.3	Effect of plant biomass	57
4.2.3.1	Catalytic performance	57

4.2.3.2	TG analysis	58
4.2.4	Effect of CP extract concentration	59
4.2.5	Effect of Pd loading	60
4.2.5.1	XRD analysis	60
4.2.5.2	AAS analysis.....	61
4.2.5.3	BET analysis	62
4.2.5.4	Catalytic performance	62
4.2.6	Effect of calcination temperature	63
4.2.6.1	FTIR analysis and TG analysis	63
4.2.6.2	XPS analysis	65
4.2.6.3	UV-Vis DRS analysis	65
4.2.6.4	TEM analysis	66
4.2.6.5	BET analysis	67
4.2.6.6	Catalytic performance	68
4.2.6.7	Stability test	70
4.2.7	Effect of H ₂ -treatment.....	71
4.2.7.1	XRD analysis	71
4.2.7.2	XPS analysis	72
4.2.7.3	UV-Vis DRS analysis	73
4.2.7.4	Catalytic performance	74
4.2.8	Effect of CeO ₂ morphology	75
4.2.8.1	TEM analysis	76
4.2.8.2	XRD analysis	76
4.2.8.3	BET analysis	77
4.2.8.4	Catalytic performance	77
4.2.9	Stability test	79
4.2.9.1	Stability of Pd/CeO ₂ /300H ₂	79
4.2.9.2	XPS analysis	80
4.3	Summary.....	81

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库